

**STUDIES ON LEAF GROWTH PARAMETERS AND SOME ELEMENTAL CONTENTS OF  
*Cucumeropsis edulis* NAUDIN INFECTED WITH *Moroccan watermelon mosaic virus-2* (MWMV)**

**UBOH, D. G., EDET, E.A., ENEH, G. D. O., UBOM, A. G. AND DANIEL, A. I.**

Department Of Science Technology  
Akwa Ibom State Polytechnic  
Ikot Ekpene, Nigeria

**ABSTRACT**

Studies on leaf growth parameters and some elemental contents of *Cucumeropsis edulis* infected with *Moroccan watermelon mosaic virus-2* (MWMV) were carried out. Test plants arranged in complete randomized designed were mechanically inoculated at 8 days except the control. Two months post inoculation, infected and healthy plants were harvested and studied for growth parameters using graphical method, weighing and linear measurements. The elemental contents of both infected and uninfected plants were evaluated using standard methods proposed by Association of Official Analytical Chemists. The results showed that MWMV caused significant reductions ( $P < 0.05$ ) in all the growth parameters of *Cucumeropsis edulis* with mean shoot height and dry leaf weight being  $52.82 \pm 6.15$  cm and  $0.40 \pm 0.31$  g in infected sample compared to the corresponding healthy values of  $75.00 \pm 13.24$  cm and  $1.23 \pm 1.00$  g obtained for the control. All the elemental contents of MWMV – infected plant were significantly ( $P < 0.05$ ) reduced with copper having the least mean value of  $3.45 \pm 0.00$  mg/100g while uninfected sample had  $5.18 \pm 0.03$  mg/100g. There is need to control the spread of this virus through phytosanitational measure, use of resistant variety and effective quarantine services. This will provide food security and poverty alleviation.

**KEYWORDS:** *Cucumeropsis edulis*, Virus and Growth.

**INTRODUCTION**

*Cucumeropsis edulis* Naudin is a species of melon belonging to the family *Cucurbitaceae*. The Yoruba, Hausa and Igbo people of Nigeria refer to the plant as “Egusi-itoo”, “Agushi” and “Ebele” respectively. The Efiks and Ibibios of Cross River and Akwa Ibom States call it “Ikpan eyong”. In tropical Africa, *Cucumeropsis edulis* is grown for food and as a source of oil (Obute *et al.*, 2007). This plant can grow up to 5-10m long, climbing by simple tendrils with stem being angular and hairy. The fruit is egg-shaped, up to about 19cm long and 8 wide and cream in colour (Obute *et al.*, 2007). The farmer sows the seeds near dead tress at the edge of garden (Egunjobi and Adebisi, 2004). In Nigeria, it is grown in the rainforest belt (Owolabi *et al.*, 2012) where the demand for the seeds particularly in towns has led to large scale planting (Adewusi *et al.*, 2000). Planting is usually between March – May at the start of the rainy season at 3-4 seeds per hole and harvested 6-8 months later. In most region, *Cucumeropsis edulis* is cultivated for cash as well as for household consumption. It is of great economic importance (Zoro Bi *et al.*, 2004). The seeds are used to prepared dough or sauce in Africa traditional socities (Ponka *et al.*, 2005).

Plant viruses are responsible for a wide range of economic losses associated with crop production. Hull (2002) defined virus as “a set of one or more nucleic acid template molecules, normally encased in protein coat, which is able to organize its own replication only within suitable host cells”. One of the plant viruses is the *Moroccan watermelon mosaic virus* (NWMV) which belongs to the genus potyvirus (Family: *Potyviridae*). This virus is characterized by flexuous rods of about 730 nm in length. It was first reported in 1972 causing severe diseases of various cucurbits in Morocco (Fischer and Lockhart, 1972). Growth and elemental alterations have been reported to occur in crops as a result of viral infection (Heil and Boston, 2002; Owolabi *et al.*, 2012). Fischer and Lockhart (1972) have remarked that *Moroccan watermelon mosaic virus* (MWMV) could be a major threat to cucurbit production. Hence, the present study was undertaken to assess the effects of *Moroccan watermelon mosaic virus* (MWMV) on growth parameters and some elemental contents of the leaf of *Cucumeropsis edulis*.

## MATERIALS AND METHODS

### Source of Seeds:

Seeds of *Cucumeropsis edulis* used in this study were sourced from Itam Main Market in Itu Local Government Area of Akwa Ibom State.

### Virus Source and Preparation of Inoculum:

The virus designated as MWMV was isolated from *Coccinia barteri* Benth by Dr. A. T. Owolabi of the Department of Botany, Faculty of Science, and University of Calabar, Nigeria. The virus was propagated and maintained on *Cucumeropsis edulis* in the greenhouse of the Department of Botany, University of Calabar, Calabar. Virus inoculum was prepared according to the procedure of Thongmearkon *et al.* (1978) by grinding virus infected *Cucumeropsis edulis* leaves with buffer (0.05 M) potassium phosphate, pH 7.5 using sterile pestle and mortar.

### Experimental Design and Inoculation of Experimental Plants:

The planting of seeds of *Cucumeropsis edulis* was carried out between March and May 2016. The plants were arranged in twenty rows of five replicates using perforated polyethylene bags each filled with 4.2 kg of treated loamy soil. The experiment was laid out in a complete randomized design with ten rows of virus inoculated plants while another ten rows served as control. Prior to inoculation (8 days after planting), the surface of the leaves were dusted with carborundum powder. Thereafter, the inoculum was applied by the conventional leaf-rub method. The leaves were rinsed with distilled water to remove superfluous inoculum so as to hasten quick infection. The plants were watered periodically and monitored for symptoms expression under the greenhouse of the Botanical Garden of the Department of Science Laboratory Technology, Akwa Ibom State Polytechnic, Ikot Osurua.

### Effect of the Virus on Shoot Height and Number of Leaves:

Sixty days post-inoculation, the effects of the virus on shoot height was determined by measuring shoot height (cm) from the base to the tip of the plants. Measurements were taken for three replicates. The leaves of inoculated and healthy plants were counted visually from each plant. Averages of triplicate determinations were considered.

### Effect of the Virus on Leaf Area:

Leaves of the same age and position on infected and healthy plants were harvested in polyethylene bags and brought to Biology Laboratory of the Department of Science Technology, Akwa Ibom State Polytechnic, Ikot Osurua. The area was traced on the graph and total area calculated based on the number of squares within the traced region (Ting, 1982).

### Effect of the Virus on Leaf Length and Width:

Infected and healthy leaves were placed on a clean specimen board before their lengths and widths were carefully measured using metre rule. Averages of triplicate measurements were taken for each plant.

### Effect of the Virus on Fresh and Dry Weights of Leaves:

Leaf fresh weight was taken for each sample by measuring using Weighing Balance. Leaf dry weight was determined by drying leaf samples at temperature of 70°C for 24 hours. Samples were dried and weighed three times using Blauscal Weighing Balance (DHG 9053A, Ocean Med. England).

### Effect of Virus on Fresh and Dry Weights of Shoots:

The harvested plants were placed in a bucket of water and the soil particles gently washed off. The shoots were cut off from the roots using scissors and then fresh weights determined and recorded. The samples were later oven-dried at 70°C for 24 hours. They were then weighed three times for both infected and healthy samples (Miyashi *et al.*, 1996).

## MINERAL ANALYSIS

The mineral analyses were conducted in the Department of Biochemistry, University of Uyo, Uyo. The infected and healthy leaf samples were oven-dried and reduced into powdery form. Digestion of the

samples followed dry digestion method after which the minerals were determined using standard procedures as described by AOAC (2005).

## STATISTICAL ANALYSIS

The Data obtained were analyzed using the student *t-test* by comparing the control with infected sample. Mean values were compared at 95% level of significant using statistical package for social science (SPSS) version 17.0.

## RESULTS

The growth parameters of *Cucumeropsis edulis* infected with *Moroccan watermelon mosaic virus-2* (MWMV) are summarized in Table 1. The results revealed that MWMV caused significant reductions ( $P < 0.05$ ) in all the growth parameters of *Cucumeropsis edulis*. The least mean value of  $0.40 \pm 0.31$  g was obtained for dry leaf weight in infected sample whereas the healthy sample had the corresponding mean value of  $1.23 \pm 1.00$  g. The virus also reduced the mean shoot height in infected sample to  $52.82 \pm 6.15$  cm whereas uninfected plant had  $75.00 \pm 13.24$  cm respectively. All the elemental contents of MWMV – infected plants were significantly ( $P < 0.05$ ) reduced (Table 2). Copper had the lowest mean value of  $3.45 \pm 0.00$  mg/100g compared to  $5.18 \pm 0.03$  mg/100g obtained for the healthy sample.

**Table 1: Growth Parameters of *Cucumeropsis edulis* Infected with *Moroccan Watermelon mosaic virus-2* (MWMV)**

Growth Parameters	Infected Sample	Health Sample
Shoot height (cm)	$52.82 \pm 6.15^*$	$75.00 \pm 13.24$
Leaf number	$7.00 \pm 0.31^*$	$12.60 \pm 1.43$
Leaf length (cm)	$5.00 \pm 0.11^*$	$9.60 \pm 1.33$
Leaf width (cm)	$7.31 \pm 0.12^*$	$8.22 \pm 0.34$
Fresh shoot weight (g)	$16.40 \pm 1.24^*$	$18.20 \pm 2.47$
Dry shoot weight (g)	$1.40 \pm 0.24^*$	$1.80 \pm 0.20$
Fresh leaf weight (g)	$7.40 \pm 0.81^*$	$10.96 \pm 1.66$
Dry leaf weight (g)	$0.40 \pm 0.31^*$	$1.23 \pm 1.00$
Leaf area (cm <sup>2</sup> )	$12.90 \pm 3.18^*$	$67.33 \pm 47.29$

Values are means  $\pm$  SEM, n = 3 replicates,  $P < 0.05$  \* significant

**Table 2: Mineral Contents of the Leaf of *Cucumeropsis edulis* Infected with *Moroccan watermelon mosaic virus-2* (MWMV)**

Minerals	Infected Sample (mg/100g)	Health Sample (mg/100g)
Potassium	$24.34 \pm 0.10^*$	$27.61 \pm 0.11$
Magnesium	$11.21 \pm 2.11^*$	$14.02 \pm 0.10$
Copper	$3.45 \pm 0.00^*$	$5.18 \pm 0.03$
Calcium	$21.22 \pm 2.30^*$	$25.11 \pm 2.10$
Iron	$6.70 \pm 0.33^*$	$9.80 \pm 0.00$
Na	$12.33 \pm 0.11^*$	$15.30 \pm 0.13$

Values are means  $\pm$  SEM, n = 3 replicates,  $P < 0.05$  \*

## DISCUSSION

This research presents the effects of *Moroccan watermelon mosaic virus-2* (MWMV) on growth parameters and some elemental contents of the leaf of *Cucumeropsis edulis*. From this study, MWMV resulted in significant reductions of all the growth parameters and elemental contents of *Cucumeropsis edulis*. This is in line with the work of Pawar *et al.* (1990) who reported reductions in shoot height, leaf weights as well as the leaf number of sorghum infected with *Sorghum ringspot virus* (SRSV). Similarly, El-DougDoug *et al.* (2005) reported that potato virus reduced the number of leaves and heights of infected plant when compared with the growth of infected sample. Attack by pathogens including viruses usually

alter plant metabolism leading to reduction in growth (Heil and Boston, 2005). Growth in plants is a complex phenomenon linked with numerous physiological processes (Owolabi *et al.*, 2012).

Elemental alterations observed in the present study are similar to the findings of Owolabi *et al.* (2010) who reported that infection of *Coccinia barteri* by MWMV caused reductions in the nutritional quality of the leafy vegetable as important dietary minerals such as magnesium, iron, calcium and vitamins A and C were significantly reduced. On the other hand, Frazer (1987) confirmed that the amounts of mineral elements in virus infected plants are usually altered.

Generally, plants make use of substantial quantities of mineral elements for their growth. They can be either macro-elements (required in relatively large amounts) or micro-elements (required only in very small amounts). Plants need potassium in large amount for photosynthesis and cambial activity whereas magnesium serves as a metabolic constituent of chlorophyll (Mehrotra and Aggarwal, 2006). The micronutrients such as iron and copper are required by plants in little amounts. Excessive supplies of these micronutrients have long been known to produce toxic effects on plants (Mehrotra and Aggarwal, 2006).

## CONCLUSION

In conclusion, MWMV-2 infection of *Cucumeropsis edulis* produced more significant ( $P < 0.05$ ) alterations on its growth and some mineral contents than as obtained for the uninfected plant.

## RECOMMENDATIONS

To secure food security and poverty alleviation, the following recommendations have become necessary:

1. Phytosanitation of all stages of plant propagation is needed.
2. Planting of resistant varieties should be encouraged.
3. More quarantine offices should be opened by government for certification of plant materials.

## ACKNOWLEDGEMENTS

The author wish to thank Mr. A. A. Umoh of the Department of Biochemistry, University of Uyo, Uyo, for his role in analyzing the mineral contents. Also, commended is the effort of Mr. Emem Mbong in statistical analysis of all the data obtained in this study.

## REFERENCES

1. Adewusi, H. G., Ladipo, R. O., Sarumi, M. B. and Vodouhe, A. A. (2000). *Agronomy in Nigeria*. In: Akoroda, M. O. (Ed), *Agronomy Re-union Day: A Book on the Theory and Practice of Agronomy as it has been in the last 33 years from July 1967 to the present in support of the Department of Agronomy*. Ibadan: University of Ibadan, Nigeria, p.241.
2. Association of Official Analytical Chemist AOAC (2005). *Official Method of Analysis*. Association of Official Analytical Chemists. Washington D.C., USA. Edited by Sidney Williams, 18th Edition, pp.96-105.
3. Egujobi, K. J. and Adebisi, A. A. (2004). *Cucumeropsis mannii*. In: Grubben G. J. H. and Denton, O. A. (Eds). *Plant Resource of Tropical Africa*. The Netherlands: Backhags Publishers, pp.235-237.
4. Fischer, H. H. and Lockhart, B. E. (1972). Serious Loses in Cucurbits Caused by Watermelon mosaic virus in Morocco. *Plant Diseases*, 58(3): 143-146.
5. Freezer, R. S. (1987). *Biochemistry of Virus Infected Plants*. Hertford: Research Study Press, p.143.
6. Hull, R. (2002). *Mathew's Plant Virology*. London: Academic Press Incorporated, p.47.
7. Miyashi, K., Sato, J. and Takahashi, N. (1996). Difference in the Effects of Dehusking during Formation of Seeds on Germination of Indica and Japaonia Rice (*Oryza sativa* L.) . *Annals of Botany*, 71(8): 569-604.
8. Obute, G. C., Ndukwu, B. C. and Chukwu, F. (2007). Targeted Mutagenesis in *Vigna unguiculate* (L.) Walp and *Cucumeropsis edulis* (Naudin) in Nigeria. *African Journal of Biotechnology*, 6(21): 246-247.
9. Owolabi, A. T., Rabenstein, F., Ehrig, F., Maiss, E. and Vetten, H. J. (2012). Strains of MWMV Isolated from *Lagenaria bresflora* and *Coccinia barteri* in Calabar, South East, Nigeria. *International Journal of Virology*, 8: 258-270.
10. Ponka, R., Fokou, E., Fotso, M. Achu, M. and Tchouanguiep, F. (2005). Methods of Preparation and the Energy, Protein and Mineral Values of Three Cameroonian Dislos: Corn Chaff, Nam Owondo/Ebobolo and Nam Ngon/Ebobolo. *African Journal of Food Agriculture, Nutrition and Development*, 5(1): 1-14.
11. Thongmearkom, P., Paschal, H. and Goodman, R. M. (1978). Yield Reduction in Soybean Infected with a Virus. *Physiopathology*, 68: 1549-1551.
12. Ting, I. P. (1982). *Plant Physiology*. Addison: Wesley Publishing Company, pp.170-188.
13. Zoro-Bi, I., Koffi, K. K. and Dje, Y. (2003). Characterization Botanique et Agronomicque de Trios Especies de Cucurbits Consommces en Sauce en Africque de I Quest: Citrullus Species; *Cucumeropsis mannii*, en Lagenaria Siceraria. *Biotechnology of Agronomy and Social Environment*, 7(3-4): 187-199.