

Study on Phytochemical, antimicrobial activity of *Phyllanthus niruri L.*

Shruti Sinha

Research Scholar, Dept. Of Biotechnology, Sunrise University, Alwar, Rajasthan.

Dr. Krishan Pal

Research Supervisor, Dept.Of Biotechnology, Sunrise University, Alwar,Rajsthan

Abstract

The plant species *Phyllanthus niruri L. roxb.*, a member of the , is a shoot holoparasite, and all the members of the genus are heterothrophic plants that present many morphological and physiological simplifications, such as an absence of cotyledons or radicles in their embryos, scale leaves without vascular tissue. Approximately 4000 species of the division Angiospermae, distributed among 270 genera and 22 families, have parasitic properties. Such plants can be classified as (i) hemiparasites, which contain chlorophyll; and holoparasites, in which chlorophyll is absent; (ii) obligatory parasites, which need to link to an adequate host immediately after germination; and facultative parasites, which are those that associate themselves to other plants after maturity; and (iii) shoot or root parasites, which attach themselves to a host according to the position their haustoria (modified roots) (Press & Phoenix, 2005; Shen et al., 2016).

Introduction

Parasitic plants are part of a group of taxonomically diverse organisms that include Angiospermae and Gymnospermae. Approximately 4000 species of the division Angiospermae, distributed among 270 genera and 22 families, have parasitic properties. Such plants can be classified as (i) hemiparasites, which contain chlorophyll; and holoparasites, in which chlorophyll is absent; (ii) obligatory parasites, which need to link to an adequate host immediately after germination; and facultative parasites, which are those that associate themselves to other plants after maturity; and (iii) shoot or root parasites, which attach themselves to a host according to the position their haustoria (modified roots) (Press & Phoenix, 2005; Shen et al., 2016).

The plant species *Phyllanthus niruri L. roxb.*, a member of the , is a shoot holoparasite, and all the members of the genus are heterothrophic plants that present many morphological and physiological simplifications, such as an absence of cotyledons or radicles in their embryos, scale leaves without vascular tissue and haustoria, all of which 2006). These plants present a significant reduction in the genes associated with the photosynthetic mechanism for evolutionary alteration linked to the chloroplast genome (Bungard, 2004)

and, therefore, do not exhibit the color green, which has led to the popular names “cipó-chumbo” (“lead-vine”) and “fios-de-ovos” (“golden-threads”) given to species of *Phyllanthus niruri* L. (Correa, 1984).

In India, *Phyllanthus niruri* L is usually associated with parasitism in ornamental plants and its occurrence in agricultural crops is unusual (Lorenzi, 2000). This species, originally from Chile, is common over the southeast and southern regions of the country, from the states of Minas Gerais to Rio Grande do Sul. They are perennial, non-chlorophyll, herbaceous, hairless plants, with filiform and delicate stems, exclusively propagated through seeds (Correa, 1984; Lorenzi, 2000). This plant is used in popular medicine as an anti-inflammatory and a diuretic remedy, for stomach and hepatic disorders, and for treating fresh wounds (Correa, 1984; Lorenzi, 2000). Phytochemical screening for the acute or chronic antiulcer properties of a freeze dried sample of the drug and the ethanolic fraction were executed by Alves et al. (1995), with positive results; however, no further studies have been found in existing literature and the purpose of this study is thus to contribute to greater knowledge on the subject, with a phytochemical characterization and evaluation of its antimicrobial activity and cytotoxicity in brine shrimp.

Material and Methods

Plant material

A sample of the *Phyllanthus niruri* L. , complete with flowers, was collected in september, in the Unn district of the city Muzaffarnagar (U.P).

Drug preparation, freeze-drying of the crude extract and fractions

The plant material was selected and was dried in an air-ventilated laboratory oven at 40 oC for 72 h. The dried plant was ground and extracted through percolation with ethanol 70% (v/v). The solution was concentrated at 45 oC under low pressure and freeze-dried. The yield of this lyophilized crude extract (CE) was 43.96% of the dried plant material. Part of the lyophilized material was used in the preparation of fractions, in accordance with the sequence of polarity of the solvent employed: 10 g of the CE was partition-fractionated with 100 mL of chloroform, shaken for 30 min and filtered to obtain a liquid fraction (CF). The same procedure was repeated with the residue using an equivalent volume of ethyl acetate (EAF), ethanol

(EF) and ethanol/water 1:1 (WEF), which were then evaporated at low pressure and the residues obtained (yields of 0.76%, 3.40%, 35.48% and 60.72%, respectively) were stored under refrigeration (Suffredini et al., 1999).

Phytochemical characterization

Thin-layer chromatography (TLC)

A 5.0 μL volume of 1.0% (m/v) CE and fractions, in addition to standard flavonoids quercetin, rutin and kaempferol (Merck®), at a concentration of 0.1% in methanol, was applied to 0.2 mm thick silica gel plates (Merck®). The following TLC system was used: ethyl acetate:methyl ethyl ketone:formic acid:water (57:27:5:10, v/v), visualized with NP/PEG as follows: 5% (v/v) ethanol NP (diphenylboric acid 2-aminoethyl ester, Sigma®) followed by 5% (v/v) ethanol PEG 4000 (polyethylene glycol 4000), being visualized under UV light at 366 nm (Wagner & Bladt, 1996).

The alkaloid extract was prepared with 2 g of the CE, shaken with 10 mL of 0.05 M sulphuric acid, then filtered. 1 mL of concentrated ammonia solution was added to this filtrate, which was then diluted to 10 mL with water, and extracted by shaking with 10 mL peroxide-free ether. The ether phase was evaporated to dryness and the subsequent residue was dissolved in methanol and tested through the following system: silica gel G (Merck®), chloroform:methanol:ammonium hydroxide (80:19:1, v/v), employing atropine sulfate 1% (m/v) as standard in a volume of 5 μL and direct visualization, after sprayed with Dragendorff reactive (Wagner & Bladt, 1996).

Flavonoid assay

The flavonoid content of the drug powder was determined according to Motta et al. (2005). A total of 2 g of the drug powder was accurately weighed, brought to a boil in about 20 mL of methanol 80% and maintained at the boil for 10 min. It was then cooled and filtered into a 50 mL volumetric flask. The drug was extracted twice and the solutions were cooled; they were then filtered and topped up to 50 mL with methanol 80%. This solution was diluted 2500-fold and 2 mL was transferred to a test tube, where 2 mL of aluminum chloride 5% (w/v) and 6 mL of methanol 80% were added. Readings were taken from a Beckman Coulter DU-640 spectrophotometer at 420 nm, after 30 min. A calibration curve was obtained using quercetin (Merck), with methanol 80% employed as a

blank. The result, expressed as a percentage of the total flavonoid content, represents an average of three readings.

Tannin assay

The amount of total tannin in the *Phyllanthus niruri* L. drug powder was determined using the colorimetric methodology approved by the European Pharmacopoeia (1984). The test was performed in triplicate and the spectrophotometer readings were carried out at a wavelength of 715 nm, after the addition of a recently prepared solution of phosphotungstic acid and topping up to 50 mL with a sodium carbonate 15% (m/v) solution. To calculate the amount of unabsorbed polyphenols, hide powder (Merck®) and pyrogallol standard (Merck®) were used.

Isolation of ethanolic fraction (EF) compounds

An EF was used for the isolation of compounds and Column chromatography, paper chromatography and NMR (Nuclear Magnetic Resonance) were used to isolate the substances (Markham & Geiger, 1996; Santos, 1998). A chromatographic glass column (60 cm x 3.5 cm of internal diameter) was filled with 35 g of Sigma® polyvinylpyrrolidone (PVPP) as a stationary phase and methanol was used as an eluant. Five grams of the EF was dissolved with a minimal amount of methanol, and this solution was introduced into the column. The fractions were gathered visually, directly from the column, accompanied by ultraviolet light (366 nm), and then successively concentrated in a Büchi® rotaevaporator. The concentrates were analyzed by TLC by using the chromatographic system: cellulose (Merck®), acetic acid 15% or 25% and UV 366 nm, after detection with diphenylboryloxy-ethylamine (Wagner & Bladt, 1996). Eighty fractions were obtained from the chromatographic column and, after selection by thin layer chromatographic profile likeness, sixteen fractions were regrouped. The regrouped fractions, 33-38 and 49-55, were subjected to descending paper chromatography. Whatman® 3 MM paper was used to isolate the substances and then, when necessary, Whatman® 3 to achieve final purification. The mobile phase employed was acetic acid at 15%. The papers were evaluated under ultra violet light (366 nm) and the bands marked, cut and extracted under reflux, using methanol as a solvent. Each band was filtered and concentrated in a Büchi® rotaevaporator. When there was a sufficient quantity of the substance, a nuclear magnetic resonance (NMR) spectrum was obtained. From the 33-38 fraction, a characteristic purple compound was isolated under 366 nm ultraviolet light, which was then called C1 (9 mg), and from the 49-55 fraction, a yellow compound was isolated under 366 nm ultraviolet light, and called C2 (12 mg). The C1 sample was analyzed with a Bruker®

Advances DPX 300 spectrometer, at 300 megahertz (MHz), whilst the C2 sample was analyzed on a DPX 500 spectrometer, at 500 MHz. The samples were dissolved in deuterated dimethyl sulfoxide (DMSO-d₆) solvent.

Biological assays

Antimicrobial activity of CE The dilution in a liquid medium (Farmacopeia Brasileira, 1988) method was used. In this study, the microorganisms used for the assay were *Staphylococcus aureus* ATCC 6538, *Pseudomonas aeruginosa* ATCC 9027 and *Escherichia coli* ATCC 10536, *Candida albicans* ATCC 10231 and *Aspergillus niger* ATCC 16404. The bacterial strains were maintained in casein soy broth (Difco), whilst the yeasts and molds were maintained in Saboraud dextrose broth (Difco). The CE was dissolved in ethanol 70% at 0.5, 1, 1.5 and 2 mg/mL concentrations. A secondary standard of chloramphenicol (2.0, 5.0 and 8.0 µg/mL), for bacteria, and amphotericin B (1, 1.5, 2, 3 and 4 µg/mL), for yeast and mold, were used as positive inhibition controls (Silva, 2001). The tubes were incubated for 24 h, or until development of the microorganisms in the control tubes, at a temperature of 30-35 °C, for bacteria, and 20-25 °C for mold. The assay was executed in triplicate. A subculture was made in the tubes that presented a precipitate but did not exhibit turbidity.

*Cytotoxicity assay (lethality of *Artemia franciscana*)*

This assay was performed according to Suffredini (2000). Initially, the potential of the CE was ascertained by using a single 1 mg/mL dose. As there was a 50%, or greater, mortality rate among the brine shrimp within 24 h, DL50 (Lethal Dose 50) was determined. The crustaceans (*Artemia franciscana*) were reared from dried eggs, which are viable for several years in this state. About 50 mg of the eggs were placed into a rectangular opaque plastic container with artificial seawater, prepared from a mixture of commercial salt, according to the manufacturer's recommendations (São Francisco Bay Brand # 65034). After 48 h, the brine shrimp that hatched were phototropically drawn to a cold light source. Ten brine shrimp were collected in a Pasteur pipette, and placed in 10 mL flasks, with 5 mg of the sample to be tested. The volume was topped up to 5 mL with seawater (concentration: 1 mg/mL). Illumination was kept constant and the survivors were counted after 24 and 48 h. For the assay to ascertain DL50, the 0.5 mg/mL, 0.1 mg/mL and 0.02 mg/mL concentrations of the CE were tested in triplicate, following the same procedure as before. DL50 was ascertained by counting the surviving brine shrimp after 24 h, and the calculation was carried out using the Taxonomy software program. The EF, WEF and extracted fraction as per Wagner & Bladt (1996) for the alkaloid compounds were also tested.

Results and Discussion

The genus *Phyllanthus niruri* L. is taxonomically controversial because it is classified as belonging to, but due to its parasitic properties, a number of schemata propose it to be a member of the monogeneric *Phyllanthus niruri* L.aceae family (Cronquist, 1981; Dahlgren, 1981). Out of the approximately 180 species that compose the genus *Phyllanthus niruri* L. (Costea et al., 2008), many are difficult to identify or differentiate, because their inherent taxonomic properties are almost entirely related to their flowers, fruits and inflorescence, whereas their vegetative parts, characterized by their very simple structures, present great uniformity (Correa, 1984; Löffler et al., 1997). The presence of flavonols derived from kaempferol and quercetin has been investigated in specimens of the Convolvulaceae family by many researchers, whilst flavonols like quercetin, kaempferol and hyperoside have been isolated in *C. micrantha*, *C. reflexa*, *C. platyloba* and *C. chinensis* (Garcia et al., 1995; Löffler et al., 1995; Yen et al., 2008). Bacchi (1993) indicated the presence of the flavonoid 3',4',5,7- tetramethyl ether of quercetin in *C. Reflexa roxb*. In this study, the thin-layer chromatographic profile (Figure 1) of the CE and the fractions obtained from the latter (WEF, EF, EAF and CF) indicates the presence of flavonoids, when revealed with diphenylboryloxy-ethylamine (Wagner & Bladt, 1996). And the assay ascertained that the drug powder consisted of 2.79% flavonoids and 2.01% tannins.




Figure 1. Chromatographic profile of extracts of *C. racemosa*. CE: lyophilized raw extract; WEF: ethanolic fraction 50%; EF: ethanolic fraction; EAF: ethyl acetate fraction; CF chloroform fraction; Rut: rutin; Querc: quercetin; canf: kaempferol. Revealed with diphenylboryloxy-ethylamine and UV light (Wagner & Bladt, 1996).

The ethanolic fraction was subjected to the chromatographic column using adsorbent polyvinylpyrrolidone, and paper chromatography was employed to purify and isolate the compounds. Two substances (**1** and **2**) were isolated and analyzed by NMR. From the data found in literature pertaining to studies of flavonoids in this genus, the proton (¹H) and carbon (¹³C) spectra of **1** (Table 1) were compared with data from literature, which indicated hyperoside (Markham & Geiger, 1996). For the compound **2**, attributions of ¹H and ¹³C NMR+HMBC spectroscopic data (Table 2), led to the conclusion that it was the 4'-methoxyquercetin flavonoid. In the antimicrobial assay, the CE presented activity with regards to the gram-positive bacteria *Staphylococcus aureus*, with a minimum inhibitory concentration of 2 mg/mL. Doses of 0.5, 1 and 1.5 mg/mL also indicated some activity, although when the subculture was made, there was growth, proving the viability of such microorganisms. This extract did not have any effect on *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans* and *Aspergillus niger*. Considering that phenolic compounds are recognized antimicrobial substances, the activity before the *Staphylococcus aureus* can be attributed to the presence of flavonoids and tannins. However, Rios et al. (1988) suggested a concentration of 100 µg/mL for plant extracts to present some antimicrobial potential and, accordingly, the value of 2 mg/mL, found in this study, would not be a promising result. The CE was effective with a single dose of 1 mg/mL in the preliminary assay performed in the brine shrimp lethality test, causing the death of 50% of the crustaceans within 24 h, and 100% within 48 h, which is enough to characterize a positive result in terms of this activity (Suffredini, 2000) and to proceed with the test to ascertain DL50. The result of DL50 was found to be 231 µg/mL. Significant values for this assay correspond to numbers lower than 1000 µg/mL (Suffredini, 2000). Therefore, the CE presented a significant result in this assay, although, when compared to the DL50 of the vincristine sulphate standard (1.2 µg/mL) obtained by Suffredini (2000), the CE value found presented a less active extract. As a result of the values obtained, new tests were carried out, with EF, WEF and with an extract prepared as per Wagner & Bladt (1996), for alkaloids. Upon analysis of the results after 24 h, the EF and WEF did not show pertinent activity for continuity and determination of

Table 1. ¹H and ¹³C NMR spectroscopic data of compound **1** (DMSO-d₆, δ ppm, 300 MHz).

No	¹ H NMR*	Encountered	¹³ C NMR (ppm)*	Encountered
2	-	-	156.3	157.1
3	O- gal	O- gal	133.8	134.3
4	-	-	177.5	178.3
5	12,61 (OH)	***	161.2	162.1
6	6,22d	6,20d	98.6	99.5
7	OH	OH	164.0	164.9
8	6,42d	6,41d	93.4	94.3
9	-	-	156.3	157.1
10	-	-	104.0	104.7
1'	-	-	121.3	121.9
2'	7,55d	7,53	115.2	116.0
3'	OH	OH	144.7	145.7
4'	OH	OH	148.5	149.3
5'	6,84d	6,83d	116.2	116.8
6'	7,68dd	7,68dd	121.8	122.8
1''	-	-	102.3	102.6
2''	-	-	71.3	72.0
3''	-	-	73.4	74.0
4''	-	-	68.0	68.7
5''	-	-	75.8	76.7
6''	-	-	60.8	60.95

*For hyperoside (Markham & Geiger, 1996). ** field not registered.

Table 2. ¹H and ¹³C NMR+HMBC spectroscopic data of compound **2** (DMSO-d₆, δ ppm, 500 MHz).

Compound 2			
No	¹ H Groups (ppm)	¹³ C NMR+HMBC (ppm)	Groups
2		150.6	— C —
3	9.51	139.8	— C — OH
4		179.9	C = O
5	12.55	164.7	— C — OH
6	6.29	102.2	— CH
7	10.89	167.9	— C — OH
8	6.58	97.6	— CH
9	-	160.1	— C —
10	-	107.0	— C —
1'	-	125.7	— C —
2'	7.79	115.7	— CH
3'	9.84	152.8	— C — OH
4'	3.94	151.4	— C — OH
5'	7.03	119.5	— CH
6'	7.85	126.0	— CH
4'	-OMe -	59.8	— OCH ₃

DL50. However, the alkaloid-directed extract ascertained by TLC (Wagner & Bladt, 1996) proved to be effective in the assay, by eliminating all the brine shrimp within 24 h. Mavlonov et al. (2008) evaluated cytotoxic activity in three species of *Phyllanthus niruri* L. (*C. europea*, *C. lupuliformis* and *C. attenuate*) in a model of murine cancer and ascertained elevated cytotoxic activity of a component of a polar fraction (aqueous extract) of *C.reflexa*. For some specimens of the genus *Phyllanthus niruri* L., alkaloids with indolic nuclei were discovered, which are considered to be potential antitumor substances (Nisa et al., 1986). Hofmann & Tschertter (Chao & Der Marderosian, 1973) were the first researchers to isolate ergoline-type alkaloids (indolic nuclei) in superior plants, since before the 1960s, this kind of alkaloid had only been found in the fungi of the genus *Claviceps*. Since the Aztec times (Mexico), the seeds of *Rivea corymbosa* and *Ipomoea violacea* (specimens of the Convolvulaceae family) have been used in religious ceremonies for hallucinogenic purposes. A wide range of ergoline-type alkaloids has been confirmed in plants of the genus *Argyreia*, as well as in others of the Convolvulaceae family, like *Rivea*, *Stictocardia* and also

Phyllanthus niruri L. (*C. monogyna* and *C. chinensis*). Pérez-Amador et al. (1996), working with four species of the genus *Phyllanthus niruri* L., after determining the chromatographic profiles of ergolinic alkaloids, glycoresins and kaurenoic glycosides (diterpenoids), which had been suggested as taxonomic markers of the Convolvulaceae family, considered them part of the Convolvulaceae family, rather than a classification apart, as suggested by Cronquist (1981) and Dahlgren (1981).

Conclusion

In addition to other species of the genus *Phyllanthus niruri* L. presented flavonoid compounds (2.79%) as the main metabolite. The presence of flavonol, reported by several authors, was also revealed in this study (4'-methoxyquercetin). Despite flavonoids being known for several biological activities, the lyophilized raw extract of the drug in question presented modest and marginally positive antimicrobial activity for the microorganism *Staphylococcus aureus*. In the lethality assay involving *Artemia franciscana*, the extract, in the form of a single 1 mg/mL dose, was able to eliminate 50% of the brine shrimp within 24 h and 100% within 48 h, resulting in a DL50 of 231 µg/mL. Although some flavonoids are known for their cytotoxic activity, the ethanolic and 50% ethanolic extracts did not prove to be effective; however, the alkaloid-directed extract was able to eliminate 100% of the brine shrimp within the 24-h period, indicating very significant activity. The cytotoxicity results obtained here, associated with the presence of alkaloids, suggested as a phytochemical marker. And found in the species of the genus *Phyllanthus niruri* L., indicate that this phytochemical group is worthy of future studies involving the species.

References

- Alves NB, Seki FR, Bacchi EM 1995. Ação antiúlcera de *Phyllanthus niruri* L. *racemosa* Martius. *Reunião de iniciação científica na FCF/USP*. São Paulo, Brazil.
- Bacchi EM 1993. Flavonoids from *Phyllanthus niruri* L. *racemosa*. *Annual congress on medicinal plant research*. Düsseldorf, Germany.
- Bungard RA 2004. Photosynthetic evolution in parasitic plants: insight from the chloroplast genome. *Bioessays* 26: 235- 247.
- Chao JM, Der Marderosian HA 1973. Identification of ergoline alkaloids in the genus *Argyreia* and related genera and their chemotaxonomic implications in the Convolvulaceae. *Phytochemistry* 12: 2435-2440.
- Correa MP 1984. *Dicionário das plantas úteis do Brasil edas exóticas cultivadas*. Rio de Janeiro: Ministry of Agriculture.

Costea M, Aiston F, Stefanovic S 2008. Species delimitation, phylogenetic relationships, and two new species in the *Phyllanthus niruri* L. *gracillima* complex (Convolvulaceae). *Botany* 86: 670-681.

Cronquist A 1981. *An integrated system of classification of flowering plants*. New York: Columbia University. Dahlgren RMT 1981. *A revised classification of the Angiosperms with comments on correlation between chemical and other characters*. In: Young DA, Seigler DS (orgs.) *Phytochemistry and angiosperm phylogeny*. New York: Praeger Publishers, p.149-204.

European Pharmacopoeia 1984. Strasbourg: Council of Europe, 2 ed.

Farmacopeia Brasileira 1988. São Paulo: Atheneu, 4a ed.

Garcia R, Erazo MS, Pena RC 1995. Flavonoids and alkaloids from *Phyllanthus niruri* L. (*Phyllanthus niruri* L. ceae). *Biochem Syst Ecol* 23: 571-572.

Löffler C, Czygan FC, Proksch P 1997. Phenolic constituents as taxonomic markers in the genus *Phyllanthus niruri* L. (*Phyllanthus niruri* L. ceae). *Biochem Syst Ecol* 25: 297-303.

Löffler C, Sahm A, Wray V, Czygan FC, Proksch P 1995. Soluble phenolic constituents from *Phyllanthus niruri* L. and *Phyllanthus niruri* L. *platyloba*. *Biochem Syst Ecol* 23: 121-128.

Lorenzi H 2000. *Plantas daninhas do Brasil*. Nova Odessa: Instituto Plantarum.

Markham KR, Geiger H 1996. ¹H nuclear magnetic resonance spectroscopy of flavonoids and their glycosides in hexadeuterodimethylsulfoxide. In: Harborne JB (org.) *The flavonoids*. London: Chapman & Hall, p.441-498.

Mavlonov GT, Ubaidullaeva KA, Kadryaeva GV, Kuznetsova NN 2008. Cytotoxic components of *Phyllanthus niruri* L.. *Chem Nat* Motta LB, Kraus JE, Salatino A, Salatino MLF 2005.

Distribution of metabolites in galled and non-galled foliar tissues of *Tibouchina pulchra*. *Biochem Syst Ecol* 33: 971-981.

Nisa M, Akbar S, Tariq M, Hussain Z 1986. Effect of *Phyllanthus niruri L. chinensis* water extract on 7,12-dimethylbenz[α] anthracene-induced skin papillomas and carcinomas in mice. *J Ethnopharmacol* 18: 21-31.

Pèrez-Amador MC, Arreola L, Márquez-Guzman J, Garcia- Argáez A 1996. Seasonal influence on the biosynthesis of ergoline alkaloids in the genus *Phyllanthus niruri L.* (Convolvulaceae). *Phyton* 59: 51-54.

Press MC, Phoenix GK 2005. Impacts of parasitic plants on natural communities. *New Phytol* 166: 737-751.

Rios JL, Recio MC, Villar A 1988. Screening methods for natural products with antimicrobial activity: a review of the literature. *J Ethnopharmacol* 23: 1127-1149.

Santos DYAC 1998. *Implicações taxonômicas dos flavonóides de Annonaceae.* São Paulo, 107 p. PhD Dissertation, Biosciences Institute of the University of São Paulo.

Shen H, Ye W, Hong L, Huang H, Wang Z, Deng X, Yang Q, Xu Z 2006. Progress in parasitic plant biology: host selection and nutrient transfer. *Plant Biol* 8: 175-185.

Silva MG 2001. *Estudo farmacognóstico de Myrciaria glomerata Berg.* São Paulo, 149 p. Master Degree Dissertation, Pharmaceutical Sciences College of the University of São Paulo.

Suffredini IB, Bacchi EM, Sertiè JAA 1999. Antiulcer action of *Microgramma squamulosa* (Kaulf) Sota. *J Ethnopharmacol* 65: 217-223.

Suffredini IB 2000. *Avaliação da atividade citotóxica, antivirale antimicrobiana de Apocynaceae amazônicas.* São Paulo, 220 p. PhD Dissertation, Pharmaceutical Sciences College of the University of São Paulo.

Wagner H, Bladt S 1996. Plant drug analysis. Berlin: Springer- Verlag.

Yen FL, Wu TH, Lin LT, Cham TM, Lin CC 2008. Concordance between antioxidant activities and flavonol contents in different extracts and fractions of *Phyllanthus niruri L. chinensis* *Food Chem* 108: 455-462.